

## Pain Assessment Tool Guidelines for use: Numeric Pain Scale

No Pain	0	1	2	3	4	5	6	7	8	9	10	Worst Pain
------------	---	---	---	---	---	---	---	---	---	---	----	---------------

### Instructions for use:

The numerical rating scale (NRS) is perhaps the most common pain assessment tool used. It is important to recognize this is an 11-point scale (0-10 not 1-10). Ask the patient whether or not they are experiencing discomfort right now. If so, ask them to describe the discomfort.

- To ascertain their understanding of the scale, ask if they would recognize:
  - a state of no pain (“0”) (or specific discomfort they describe)
  - and the worst pain (or specific discomfort) imaginable (“10”).
- Have the patient rate the intensity of their pain/discomfort “right now” verbally with a number or by pointing to the number that represents their pain intensity
- This process can be repeated with similar questions about emotional distress

### Scoring instructions:

Record the corresponding numerical value based on the patient’s response.

### Populations for use:

The NRS is a widely used tool that is appropriate for adults, and may be used in the mildly cognitively impaired adult. Patients report lower pain scores when they are educated about the proper use of this scale.

### Validity and reliability:

The validity of the NRS has been well established and it has been shown to be easy to use.

### References:

- Gracely RH, Wolskee PJ. (1983). Semantic functional measurement of pain: Integrating perception and language. *Pain, 15* :389-398.
- Herr, K.A. & Garand, L. (2001). Assessment and measurement of pain in older adults. *Clinic of Geriatric Medicine, 17*, 457-478.
- Houde RW. (1982). Methods for measuring clinical pain in humans. *ACTA Anaesthesiologica Scandinavica, 74* (Suppl) :25-29.
- Jensen MP, Karoly P, Braver S. (1986). The measurement of clinical pain intensity: a comparison of six methods. *Pain, 27* :117-126.
- Marco CA, Marco AP, Plewa MC, et al. (2006). The verbal numeric pain scale: effects of patient education on self-reports of pain. *Academic Emergency Medicine, 13*:853-859.
- Sriwatanakul K, Kelvie W. Lasagna L. (1982). The quantification of pain: An analysis of words used to describe pain and analgesia in clinical trials. *Clinical Pharmacology and Therapeutics, 32* :143-148.