Pain Relief Connection

The Pain Information Newsletter Provided by MGH Cares About Pain Relief,

a program of Patient Care Services



Archived issues are available at http://www.MassGeneral.org/PainRelief

Volume 9, No. 5			May 2010
In this issue:			
In the News	Page 1	Pain Resources on the Web	Page 2
Journal Watch	Page 1-2	Pain Education Opportunities	Page 2
CAM	Page 2	MGH Pain Calendar & Resources	Page 2

In the News

- FDA Launched a new initiative to <u>Reduce Infusion Pump Risks</u> based on reports of over 56,000 adverse events and more than 500 deaths associated with the use of infusion pumps despite 87 pump recalls.
- Dozens of over-the-counter drugs made by McNeil for infants and children, including <u>Tylenol, Motrin,</u> <u>Zyrtec, & Benadryl products cave been recalled</u> due too higher than listed drug concentrations and impurities. This follows a recall of <u>Arthritis Strength Tylenol</u> earlier this year for similar reasons.
- New Palliative Sedation guidelines delineate proper use at end of life when intolerable pain persists.
- The DEA will allow <u>electronic prescribing of controlled substance</u> effective June 1, 2010 with strings attached.

Journal Watch All items are accessible via MGH computers/library; MGHers can obtain articles from journals to which Treadwell Library does not subscribe through the <u>Treadwell home page</u>, go to 'Order Articles' under the 'Quick Picks' banner on the left.

- Fine P, Herr K, Titler M, The <u>cancer pain practice index</u>: A measure of evidence-based practice adherence for cancer pain management in older adults in hospice care. *J Pain Symptom Manage*. 2010 May;39(5):791-802. Describes a tool developed to evaluate nurse and physician adherence to pain management guidelines.
- Blinderman CD. (2010). <u>Opioids, iatrogenic harm and disclosure</u> of medical error. *J Pain Symptom Manage,* 39:309-13. A case review explores possible iatrogenic harm and asks "Is it a medical error at the end-of-life?"
- Clay F, Newstead SV, Watson WL, et al (2010). Bio-psychosocial <u>determinants of persistent pain</u> six months after non-life-threatening acute orthopedic trauma. *J Pain*, *11* (5) :420-430. High initial pain, blaming others for the injury & psychological distress are independent predictors persistent pain, its intensity & related disability.
- Claxton RN, Blackhall L, Weisbord SD, et al. (2010). <u>Undertreatment of symptoms</u> in patients on maintenance hemodialysis. *Jo of Pain & Symptom Manage, 39* :211-8. Only 45% of patients with bone/joint pain reported receiving an analgesic medication. PCP's are more likely to treat distress with analgesics than specialists.
- Nieminen TH, Hagelberg NM, Saari TI, et al. (2010) <u>Grapefruit juice enhances the exposure to oral oxycodone</u>. *Basic Clin Pharmacol Toxicol*. [epub ahead of print] Grapefruit juice does not enhance analgesic effect (despite increased in blood levels) but may increase side effects of oxycodone due to altered liver function.
- Twycross A. &Dowden SJ. (2009). Status of <u>pediatric nurses' knowledge about pain</u>. *Pediatric Pain Letter*, 11 (3): 17-21. accessed @ <u>www.childpain.org/ppl</u>. Nursing education, experience and professional activities (e.g. participation with organizations or committees) contribute to knowledge about pediatric pain management.
- Anand KJ, Willson DF, Berger J, et al. <u>Tolerance and withdrawal from prolonged opioid use</u> in critically ill children. *Pediatrics*, 2010;125(5):e1208-e1225 Opioid tolerance is more profound in younger children, during critical illness, and when short-acting opioids are used for longer periods. Treatment options are reviewed.
- Weng YM, Chang YC, Lin YJ. (2010). <u>Triage pain scales cannot predict analgesia</u> provision to pediatric patients with long-bone fracture. *Am J Emerg Med*, 28:412-7. Delays and ineffective analgesia are common.
- Reza FM, Zahra F, Esmaeel F, et al. (2010). Preemptive analgesic effect of ketamine in patients undergoing elective cesarean section. *Clin J Pain*, 26 :223-226. <u>Ketamine is not effective before a cesarean section</u>.

MGH Cares About Pain Relief

Massachusetts General Hospital PainRelief@Partners.org • http://www.MassGeneral.org/PainRelief

To be added to or removed from the Pain Relief Connection mailing list, send an email to pmarnstein@partners.org

Journal Watch (continued)

- Huijnen IP, Verbunt JA, Peters ML, et al. (2010). Do <u>depression and pain intensity</u> interfere with physical activity in daily life in patients with Chronic Low Back Pain? *Pain* [ePub ahead of print]. The extent of depression with CLBP better predicts functioning (both objective & subjective measures) more than pain level.
- Eisenach JC, Curry R, Rauck R, et al. (2010) Role of <u>Spinal Cyclooxygenase</u> in Human Postoperative and Chronic Pain. *Anesthesiology*, *112*(5):1225-1233. Based on the lack of effect from intrathecal ketorolac, it was concluded that spinal cylcooxygenase activity does not contribute to chronic or postoperative pain.
- Tang NK, Goodchild CE, Hester J, et al. (2010). <u>Mental defeat</u> is linked to interference, distress and disability in chronic pain. *Pain*, *149* (3):547-54 2010. Pain patients who feel mentally defeated reported a greater degree of pain, its interference with daily life, distress and disability. This concept joins with other potentially destructive psychosocial factors like anxiety, depression, helplessness and catastrophizing linked to disability.
- Moore RA, Straube S, Paine J, et al. (2010). <u>Fibromyalgia: Moderate and substantial pain intensity reduction</u> predicts improvement in other outcomes and substantial quality of life gain. *Pain, 149* (2):360-364. Often people with fibromyalgia continue with ineffective therapies. Pain reduction is a simple and effective predictor of which patients should continue treatment, and which should discontinue and try an alternative therapy

CAM

- The <u>pain relieving properties of ginger</u> supplementation (raw or heated) cuts exercise-induced pain by 25%.
- Interpersonal and therapeutic skills may be more important in acupuncture than training of the acupuncturist.
- <u>Slow breathing reduced ratings of pain intensity and unpleasantness</u>, particularly for moderate vs. mild pain.

Pain Resources on the Web

- <u>GeriatricPain.org</u> has many evidence-based tools that help nurses to better manage pain in older adults.
- The <u>American Chronic Pain Association</u> has a revised, easier to navigate website for people living with pain.

Pain-Related Education Opportunities

- Wed-Thu June 16-17th the Art & Science of Palliative Nursing held at Northeastern University, Boston, MA
- Tue-Sat June 22-26th Principles and Practice of Pain Medicine CME course of Harvard Boston, MA
- Thu-Sun June 24-27th Annual (52nd) Scientific Meeting of <u>American Headache Society</u> LA, CA

MGH Pain Calendar

- Chronic Pain Rounds are held Mondays at 12:00N in the Ether Dome. Mailing: ttoland@partners.org
- Palliative Care Grand Rounds held Wednesdays at 8:00am in the Ether Dome. Mail: nalawless@partners.org

MGH Pain Resources:
PainRelief web site: http://www.massgeneral.org/painrelief/
Previous issues of *Pain Relief Connection*: MGH Center for Translational Pain Research: MGH Center for Translational Pain Research

MGH PCS educational offerings
http://www2.massgeneral.org/pcs/The_Institute_for_Patient_Care/EDU/Educational_Offering_Events_Calendar.pdf

Treadwell Library (Magic): http://magic.mgh.harvard.edu/

MGH Pain Medicine: http://magic.mgh.harvard.edu/

MGH Formulary (includes patient teaching handouts in 16 languages): http://www.crlonline.com/crlsql/servlet/crlonline

Partners Handbook: http://is.partners.org/handbook/

Primary Care Office InSite (PCOI) (Clinician and patient information): http://intranet.massgeneral.org/pcs/Pain/index.asp

MGH Cares About Pain Relief

Massachusetts General Hospital

PainRelief@Partners.org • http://www.MassGeneral.org/PainRelief

To be added to or removed from the Pain Relief Connection mailing list, send an email to pmarnstein@partners.org