



Volume 11, No.9

MGH CARES ABOUT PAIN RELIEF

September 2012

PAIN RELIEF CONNECTION THE PAIN INFORMATION NEWSLETTER

Provided by MGH Cares About Pain Relief, a program of MGH Patient Care Services

Inside this issue:

n the News	Page 1	Pain Resources on the Web	Page 2
Journal Watch	Page 1 - 2	Pain Education Opportunities	•
CAM	Page 2	MGH Pain Calendar & Resources	•

In the News

- FDA summit concluded we can't predict which chronic pain patients will be helped, hurt, or receive no benefit from chronic opioid therapy.
- The International Association for the Study of Pain has released its <u>Interprofessional Education Curriculum</u> for health professional training
- National drug take back day is Sept 29th. Rid your & patients' medicine cabinet of unwanted or expired drugs for safe & proper disposal.
- FDA has information for the public and patient advocates on safe use of analgesics, including <u>Acetaminophen</u> and <u>NSAIDs</u> and <u>Opioids</u>

Journal Watch

All items are accessible via MGH computers/library. MGHers can obtain other articles from through the Treadwell home page

- Pizzi LT, Toner R, Foley K, et al. (2012). Relationship between potential opioid-related adverse effects and hospital length of stay in patients receiving opioids after orthopedic surgery. *Pharmacotherapy*, 32(6):502-14. After orthopedic surgery opioid-related constipation or confusion added more than a day to length of stay. Nausea and vomiting are more common but impact length of stay less.
- Haroutiunian S, Donaldson G, Yu J, et al. (2012). Development and validation of shortened, restructured Treatment Outcomes in Pain Survey instrument (the S-TOPS) for assessment of individual pain patients' health-related quality of life. *Pain*, *153* (8): 1593-601. A new valid scale that measures aspects of quality of life for pain patients, including pain intensity, functioning & patient's rating of improvement.
- Rustøen T, Valeberg BT, Kolstad E, et al. (2012) The Pro-Self® pain control program improves patients knowledge of cancer pain management. *J Pain Symptom Manage*, 44(3):321-30. A <u>nurse coached intervention helps bone cancer pain</u> patients' outlook & function.
- Logan DE, Conroy C, Sieberg CB, et al. (2012). Changes in <u>willingness to self-manage pain among children</u> and adolescents and their parents enrolled in an intensive interdisciplinary pediatric pain treatment program. *Pain*, *153* (9):1863-70. Helping children and parents self-manage pain is linked to a reduction in pain and the fear of it; functional improvement, and the use of more adaptive coping skills.
- Soto E, Stewart DR, Mannes AJ, et a.. (2012). Oral ketamine in the palliative care setting: a review of the literature and case report of a patient with neurofibromatosis type 1 and glomus tumor-associated complex regional pain syndrome. *Am J Hosp Palliat Care*, 29 (4):308-17. Excellent review and dosing guidance for oral ketamine, increasingly used as a pain management tool, especially for outpatients.
- Echevarria G, Elgueta F, Fierro C, et al. (2011). Nitrous oxide (N2O) reduces postoperative opioid-induced hyperalgesia after remifentanil-propofol anaesthesia in humans. *Br J Anaesth*, *107*(6):959-65. Pain thresholds are not lowered as much 12-18 hours after surgery. Remifentanyl may lower thresholds and patients receiving intraoperative Nitrous Oxide, may protect against this effect.

MGH Cares About Pain Relief
Massachusetts General Hospital
PainRelief@partners.org ● http://www.mghpcs.org/painrelief

Journal Watch (continued) [MGHers can obtain articles through the <u>Treadwell home page</u>]

- Pinto PR, McIntyre T, Almeida A, et al. (2012). The mediating role of pain catastrophizing in the <u>relationship between presurgical anxiety</u> and acute postsurgical pain after hysterectomy. *Pain*, *153*(1):218-26. Women with high preoperative pain were 2.5 times more likely to have high postoperative pain at 2 days after surgery. Age, anxiety, unhelpful thoughts & comorbid conditionals also predicted more pain.
- Clarke H, Bonin RP, Orser BA, et al. (2012). The prevention of chronic postsurgical pain using gabapentin and pregabalin: A combined systematic review and meta-analysis. *Anesth Analg*,15(2):428-42. <u>Preemptive gabapentin or pregabalin cuts risk of prolonged pain.</u>
- Caraceni A, Hanks G, Kaasa S, et al. (2012) Use of opioid analgesics in the treatment of cancer pain: evidence-based recommendations
 from the EAPC. Lancet Oncology. 13(2):e58-68. A limited research base with publication bias makes GRADE evidence based guidelines
 challenging. Biggest changes recommended are Step II should remove codeine & tramadol in favor of low-dose oxycodone or morphine.
- Macfarlane GJ, Beasley M, Jones EA, et al. (2012). The <u>prevalence and management of low back pain across adulthood</u>: results from a population-based cross-sectional study (the MUSICIAN study). *Pain*, 153(1):27-32. Older adults have double the prevalence of disabling low back pain than those under age 40. Having previously failed exercise-based therapy, elders are often treated with medicine alone.

Pain Resources on the Web

- Excellent online program by Dr. Roger Chou entitled, "Clinical Guidelines for Opioid Use in Chronic Noncancer Pain".
- A free log in gets you the latest National Comprehensive Cancer Network (NCCN) 2012 guidelines on cancer pain management
- GeriatricPain.org is a website filled with useful tools and information on pain in the older adult, with a focus on those in nursing homes.

CAM (Complementary and Alternative Medicine)

- In the largest study of its kind, the nutritional supplement <u>L-carnitine failed to reduce pain</u>, depression or fatigue in cancer patients.
- Warming infants during immunizations may be better than sucrose tasting (e.g. Sweet Ease) or pacifier sucking.
- Modest benefits of acupuncture are not related to placebo effect or attitude. When exposed to (ice or capsaicin) induced pain.

Pain-Related Education Opportunities

- Sat September 29th The Art and Science of Pediatric Pain and Management conference at Boston Children's Hospital see brochure
- Tue-Sun Nov 15-18, 2012 American Society of Regional Anesthesia meeting focuses on invasive & non-invasive therapy Miami, Florida

MGH Pain Calendar

- Fri. October 5th Need-to-know Basics of Pain Management: Founders 325 at 8-9am email for info repeated November 9th
- Mon Oct 29th Gaining Control of Complex Pain. O'Keeffe Auditorium 8am 4pm email for info
- Mon November 19th 11am 3pm, Beyond the Basics of Pain Management Founders 325 email for info
- Chronic Pain Rounds occur weekly on Mondays at 12:00N Mail: email <u>Tina Toland f</u>or details
- Palliative Care Grand Rounds occur weekly on Wednesdays at 8:00am. email: Margaret Spinale

MGH Pain Resources

The Patient Education Television: Dial 4-5212 from patient's phone then order: #279 for Chronic Pain; #280 for Cancer Pain; #281 for Communicating Pain; #282 for Prescription and Non-prescription Pain Medications; @3283 for Postoperative Pain Excellence Every Day Pain Portal Page: http://www.mghpcs.org/eed portal/EED pain.asp

The MGH Center for Translational Pain Research: http://www.massgeneral.org/painresearch

MGH Pain Medicine: http://www2.massgeneral.org/anesthesia/index.aspx?page=clinical_services_pain&subpage=pain

MGH Palliative Care: http://www.massgeneral.org/palliativecare/

MGH Formulary (includes patient teaching handouts in 16 languages): http://www.crlonline.com/crlsql/servlet/crlonline
Intranet site for MGH use to locate pain assessment tools and policies: http://intranet.massgeneral.org/pcs/Pain/index.asp

MGH Cares About Pain Relief
Massachusetts General Hospital
PainRelief@partners.org ● http://www.mghpcs.org/painrelief