

Volume 19 No. 6

MGH CARES ABOUT PAIN RELIEF

June 2020

PAIN RELIEF CONNECTION THE PAIN INFORMATION NEWSLETTER

Provided by MGH Cares About Pain Relief, a program of MGH Patient Care Services

Inside this issue:

In the News	.Page 1	
Journal Watch	.Page 1-2	
CIH		

Pain Resources on the Web......Page 2 Pain Education Opportunities.....Page 2 MGH Pain Calendar & Resources.....Page 2

In the News

- Massachusetts statistics show 94% of opioid deaths in 2019 involved illicit fentanyl; while other opioid deaths declined slightly.
- FDA has granted Orphan Drug designation to begin trials of a proprietary topical amitriptyline to treat erythromelalgia.
- Symptoms of heart attack or stroke may signal COVID-19 infection, whose late effects may resemble fibromyalgia.
- Cebranopadol is a novel, centrally acting, first-in-class analgesic with a unique mechanism of action in phase III trials.
- A CDC stakeholder engagement meeting to discuss pain (acute & chronic) will be online Wednesday July 22, from 10:00am-1:00pm.
- FDA recalls injectable Cannabidiol + Curcumin whose safety & efficacy for pain has not been established.
- Australians roll out a number of regulatory changes to minimize prescription opioid harms while maintaining access when needed for pain.

Journal Watch [MGHers can obtain articles through the Treadwell home page]

- Robson EK, Hodder RK, Kamper SJ, et al. Effectiveness of <u>weight-loss interventions for reducing pain</u> and disability in people with common musculoskeletal disorders: a systematic review with meta-analysis. *J Orthop Sports Phys Ther*. 2020 Jun;50(6):319-333. Weight-loss interventions may have a small effect on pain & disability in people with knee or hip osteoarthritis, but not back pain.
- Muzaale AD, Daubresse M, Bae S, et al. <u>Benzodiazepines, co-dispensed opioids, and mortality among patients initiating long-term incenter hemodialysis.</u> Clin J Am Soc Nephrol. June 2020, 15 (6) 794-804. Benzodiazepines are often prescribed to hemodialysis patients & is associated with a 20% higher risk of death with short-acting agents and a 60% higher mortality rate when co-prescribed with an opioid.
- Hayes CJ, Krebs EE, et al. Impact of opioid dose escalation on pain intensity a retrospective cohort study. *Pain*. 2020 May;161(5):979-988. VA databases suggest for chronic noncancer pain, a <u>stable opioid dose</u>, is better than escalating opioid doses over a 6-month period.
- Friedman BW, Ochoa LA, Naeem F, et al. Opioid use during the six months after an emergency department visit for acute pain: a prospective cohort study. *Ann Emerg Med*. 2020 May;75(5):578-586. Although 21% of opioid-naïve patients filled up to 2 opioid prescriptions after being prescribed an opioid in the ED, only 1% had persistent opioid use for ongoing moderate-severe pain at 6 months.
- Goyal MK, Johnson TJ, Chamberlain JM, et al. Racial and ethnic differences in emergency department pain management of children with fractures. *Pediatrics* May 2020, 145 (5) e20193370 Minority children with long bone fractures often get less opioids / adequate pain control.
- Chino F, Kamal A, Chino J. Incidence of opioid-associated deaths in cancer survivors in the United States, 2006-2016: a population study of the opioid epidemic. *JAMA Oncol.* ePub online May 7, 2020. Cancer patients die from opioids at 1/10 the rate of the general population.
- Al-Ghabeesh SH, Bashayreh IH, Saifan AR, et al. <u>Barriers to effective pain management in cancer patients</u> from the perspective of
 patients and family caregivers: a qualitative study. *Pain Manage Nursing* 2020 Jun 21(3), p238-44. Regulatory factors, knowledge deficit,
 and the use of religious and cultural strategies to cope with pain were major barriers to effective cancer pain management.

MGH Cares About Pain Relief
Massachusetts General Hospital

<u>Previous Newsletters</u>

Journal Watch (continued) [MGHers can obtain articles through the Treadwell home page]

- Bevil KM, Klesius LL, Chambers T, et al. <u>Educating perioperative purses about local anesthetic systemic toxicity</u> (LAST) using high-fidelity simulation. *Pain Manage Nursing* 2020 Jun 21(3), p265-70. Teaching perioperative nurses about LAST using a simulation where the patient progressed from mild to life-threatening effects were found to significantly improved knowledge that was retained for 3 months
- Donnelly TJ, Palermo TM, Newton-John TRO. Parent cognitive, behavioural, and affective factors and their relation to child pain and functioning in pediatric chronic pain: a systematic review and meta-analysis. *Pain*. 2020 Jul;161(7):1401-1419 Parental mood, thoughts & protective behaviors linked to poorer school functioning & disability among children with chronic pain.
- Searle S, Muse D, Paluch E, et al. Efficacy and safety of single and multiple doses of a fixed-dose combination of <u>ibuprofen and</u> <u>acetaminophen in the treatment of postsurgical dental pain</u>: Results from two studies. Clin J Pain. 2020 Jul;36(7):495-504. Fixed-doses of ibuprofen (IBU) 250mg + acetaminophen (APAP) 500mg were better than either IBU or APAP alone after ≥ 3 molar extractions.
- Fredrikson KM, Fasolino T. Pharmacogenetic testing: the ethics of implementing in clinical practice for chronic pain patients. *J Pain Palliat Care Pharm*. 2020;34(2):69-76. Is it our ethical duty to do <u>use pharmacogenetics to promote safe</u>, effective pain treatments, not trial & error?
- Depelteau A, et al. Chronic pain and frequent use of emergency department: A systematic review. *Am J Emerg Med*. 2020, Feb;38(2):358-363. Chronic pain patients are high utilizers of the emergency room, with higher impact (disabling) chronic pain had more frequent visits.
- Ojo OA, Mehdiratta JE, Gamez BH, et al. Comparison of Programmed Intermittent Epidural Boluses (PIEB) with Continuous Epidural Infusion
 (CEI) for the maintenance of labor analgesia: a double-blind RCT study. Anesth Analg. 2020;130:426-435. Epidurals in women during labor
 with PIEB had less motor bock than those with a continuous epidural infusion. Both had similar patient controlled epidural doses received.
- Wei J, Lane NE, Bolster MB, et al. Association of tramadol use with risk of hip fracture. *J Bone Miner Res.* 2020 Apr;35(4):631-40. Among adults over age 50 in an 18 year study, those placed on tramadol had a higher risk of hip fracture than those on codeine or NSAIDs.

Pain Resources on the Web:

- Physiologic evidence now shows African Americans experience pain as more intense unpleasant and disabling than whites
- <u>European Pain Federation position papers</u> include timely topics, including a new one on opioid prescribing to prevent a US-type opioid crisis.
- The Emergency Medical Services for Children has a <u>Pediatric Pain Management Toolkit</u> for professionals, caregivers & organizations.

Complementary Integrative Health

- Some psychologic therapies cut chronic pain intensity in 4 sessions, including hypnosis, cognitive therapy & pain education.
- Multiple clinical trials show kinesio-taping effectively reduces pain & improves knee functioning with arthritis.
- A single shot of vitamin D lowered neuropathic pain and improved balance in diabetic patients with low vitamin D levels.
- Singing as a form of music therapy helps people with chronic pain develop a sense of physical, psychological & social well-being

Pain-Related Education Opportunities

- Jul 28, 8:00am <u>Educating Healthcare Professionals about Pain Prevention</u>. An international global year for pain prevention webinar.
- FDA-approved Risk Evaluation and Mitigation Strategies (<u>REMS</u>) for opioids updated online training & resources
- NEJM Knowledge+ Pain Management best practices and Safe Opioid Prescribing Principles with CME credits
- Scope of Pain A 6-part series on effectively manage acute &/or chronic pain, with a judicious and safer use of opioid analgesics.

MGH Pain Calendar Postponed until the fall

MGH Pain Resources

Excellence Every Day Pain Portal Page: http://www.mghpcs.org/EED/Pain/default.shtml

MGH Pain Management Center Newsletter: https://conta.cc/2Hk79qV

The MGH Center for Translational Pain Research: https://www.massgeneral.org/TranslationalPainResearch/

MGH Palliative Care: http://www.massgeneral.org/palliativecare

MGH Formulary (includes patient teaching handouts in 16 languages): http://www.crlonline.com/crlsql/servlet/crlonline

MGH Cares About Pain Relief
Massachusetts General Hospital

Previous Newsletters

To be added to or removed from the Pain Relief Connection mailing list, or other inquiries, send an email to Paul Amstern