AMERICAN JOURNAL OF PREVENTIVE MEDICINE

Am J Prev Med. 2020 Apr 27

doi: 10.1016/j.amepre.2020.04.003 [Epub ahead of print]

PMCID: PMC7183932

Elsevier

Disparities in the Population at Risk of Severe Illness From COVID-19 by Race/Ethnicity and Income

Matthew Raifman, MPP^{1, □} and Julia Raifman, ScD²

Author information Copyright and License information Disclaimer

Associated Data

Supplementary Materials

Go to:

INTRODUCTION

Identifying those at heightened risk of severe illness from novel coronavirus disease 2019 (COVID-19) is essential for modeling disease, designing return-to-work criteria, allocating economic assistance, advancing health equity, and limiting morbidity and mortality. The U.S. Centers for Disease Control and Prevention has identified criteria associated with risk of severe complications from COVID-19 infection (Appendix Table 1).¹ Structural inequities have shaped racial, ethnic, and income disparities for many of these criteria. To date, there has been limited analysis of the proportion of the population at risk in the U.S. based on these criteria,² or risk factors by race/ethnicity or income. Preliminary national data on cases by race/ethnicity suggest that disparities in hospitalization are already developing.³ Quantifying disparities in risk is important for allocating resources to prevent, identify, and treat COVID-19-related severe illness and limit diverging outcomes for already vulnerable subgroups.

Go to:

METHODS

The authors used data from the 2018 Behavioral Risk Factor Surveillance System, a nationally representative survey of >400,000 adults. This study estimated the proportion of adults that have at least one of the Centers for Disease Control and Prevention criteria for risk of severe illness from COVID-19 (hereafter "higher risk") (Appendix Table 1) by age group, and by race/ethnicity and household income (<\$25,000 or not), accounting for comorbidities. In each age group, the proportion with more than one risk factor was also estimated, recognizing the potential for compounding risk. Prevalence ratios (PRs) and 95% CIs were estimated for

race/ethnicity and income group using Poisson regression models with robust SEs. The authors excluded those who responded *don't know/not sure* or *refused*, or had missing data for any risk factor except pregnancy (Appendix Tables 1 and 2).

Go to:

RESULTS

There were 387,304 respondents for the overall and race/ethnicity analyses, and 336,861 respondents for the income analysis (Appendix Table 3).

Approximately 97 million Americans (43% of adults) were at higher risk of illness from COVID-19. Among those aged <65 years, 28% were higher risk, including 18% of the youngest age group (18–24 years).

Racial/ethnic disparities in risk were found across age groups (<u>Figure 1</u> A). Among those aged <65 years, 33% of black (PR=1.22, 95% CI=1.18, 1.26) and 42% of American Indian (PR=1.53, 95% CI=1.41, 1.66) adults were at higher risk relative to 27% of white adults (Appendix Table 4).




Figure 1

Disparities in prevalence of CDC risk factors for severe illness from COVID-19 by race/ethnicity and income.

W, white; H, Hispanic; B, black; AI, American Indian; A; Asian

More than 40 million Americans (18%) had two or more risk factors. Among those aged <65 years, 11% of black (PR=1.27, 95% CI=1.19, 1.36) and 18% of American Indian (PR=2.15, 95% CI=1.84, 2.50) people had multiple risk factors relative to 8% of white people (Appendix Tables 5 and 8). Among those aged \geq 65 years, 69% of American Indian (PR=1.28, 95% CI=1.18, 1.38) and 61% of black (PR=1.12, 95% CI=1.08, 1.17) people had an additional risk factor(s) beyond age relative to 54% of white adults (Appendix Table 6).

At least 25 million Americans living in low-income households were higher risk, and there were also disparities across age groups (Figure 1B). Among those aged <65 years, 40% of low-income people were higher risk (PR=1.63, 95% CI=1.59, 1.67) relative to 24% of those with higher income (Appendix Table 7). There were also income disparities in multiple risk factors among those aged <65 years (16% vs 6%, PR=2.51, 95% CI=2.38, 2.63) (Appendix Table 8) and >65 years (63% vs 52%, PR=1.21, 95% CI=1.18, 1.24) (Appendix Table 9).

Go to:

DISCUSSION

People who are black, American Indian, or live in low-income households are more likely to have conditions associated with increased risk of illness from COVID-19 relative to those who are white or higher income, respectively. These inequities in risk are compounded by structural disparities in access to medical insurance,⁴ wealth, and income volatility.⁵

Structural inequities also contribute to heightened exposure to COVID-19. Minorities and people living in low-income households are more likely to work in industries that have remained open during non-essential business closures.⁶ They are also more likely to live in crowded conditions² and multigenerational households that may elevate exposure and limit options for quarantining family members.

It is vital that these race/ethnicity and income disparities in risk be considered in physical distancing policies and other protective measures, particularly for those who work in essential industries. Rationing resources based on comorbidities may exacerbate inequities, whereas prioritizing vaccine delivery on risk may reduce deaths and disparities.

Data availability in the Behavioral Risk Factor Surveillance System is a limiting factor; the data do not include all risk criteria and only capture respondents who were aware of their condition. These estimates represent a lower bound of adults at risk of severe illness. It is possible that risk is not uniform and those with multiple factors may be at higher risk. This would be further evidence of race/ethnicity and income disparities.

Go to:

ACKNOWLEDGMENTS

We would like to acknowledge Professors Jonathan Levy, ScD and Patrick Kinney, ScD of the Department of Environmental Health at the Boston University School of Public Health, and Lorraine Dean, ScD of the Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, for their review of the manuscript and guidance. They received no compensation for this role. The research presented in this paper is that of the authors and does not reflect the official policy of the NIH. This research was supported by training grant number 5T32ES014562-13 from the NIH National Institute of Environmental Health Sciences and grant number K01 MH116817 from the NIH National Institute of Mental Health.

Both authors contributed equally to all aspects of the manuscript.

No financial disclosures were reported by the authors of this paper.

Footnotes

Supplemental materials associated with this article can be found in the online version at <u>https://doi.org/10.1016/j.amepre.2020.04.003</u>.

<u>Go to:</u>

Appendix.SUPPLEMENTAL MATERIALClick here to view.(520K, pdf)Image, application 1

<u>Go to:</u>

REFERENCES

1. CDC. People who are at higher risk for severe illness. <u>www.cdc.gov/coronavirus/2019-ncov/specific-groups/people-at-higher-risk.html</u>. Accessed March 28, 2020.

2. The Henry J. Kaiser Family Foundation. How Many Adults Are at Risk of Serious Illness If Infected with Coronavirus?<u>www.kff.org/global-health-policy/issue-brief/how-many-</u>adults-are-at-risk-of-serious-illness-if-infected-with-coronavirus/. Accessed March 28, 2020.

3. Garg S, Kim L, Whitaker M. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019 — COVID-NET, 14 states, March 1–30, 2020. MMWR Morb Mortal Wkly Rep. 2020;69(15):458–464. doi: 10.15585/mmwr.mm6915e3. [PubMed] [CrossRef] [Google Scholar]

4. Griffith K, Evans L, Bor J. The Affordable Care Act reduced socioeconomic disparities in health care access. Health Aff (Millwood) 2017;36(8):1503–1510. doi: 10.1377/hlthaff.2017.0083. [CrossRef] [Google Scholar]

5. The Pew Charitable Trusts. How Income Volatility Interacts With American Families' Financial Security. <u>www.pewtrusts.org/en/research-and-analysis/issue-briefs/2017/03/how-income-volatility-interacts-with-american-families-financial-security</u>. Accessed March 28, 2020.

6. U.S. Bureau of Labor Statistics. BLS Reports: A profile of the working poor,
2016. <u>www.bls.gov/opub/reports/working-poor/2016/home.htm</u>. Accessed March 31, 2020.

7. Adamkiewicz G, Zota AR, Fabian MP. Moving environmental justice indoors: understanding structural influences on residential exposure patterns in low-income communities. Am J Public Health. 2011;101(suppl 1):S238–S245. doi: 10.2105/ajph.2011.300119. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

<u>Go to:</u>